

Effect of Autogenic Training and Meditation on Mental Stress Management among Judo Players

Dr Adengada A Kushalappa* Dr S K Manjunatha**

*Physical Education Director Cauvery College Kodagu

**Assistant Professor Department of Physical Education, College of Horticultural, Mudigere.

(Received 12 November 2015 – Accepted & Published 26 December 2015)

Abstract

Many people are unhappy in the world, dissatisfied and yet not sure what is lacking in their lives. The basic reason for this unhappiness is our attachment to the material plane of existence. Once we gain a glimpse of higher spheres of consciousness, then our unhappiness and discontent automatically fade away. Man has made tremendous advances in almost every walk of the life. In this study the variables are independent variable: Autogenic Training and Meditation. Dependent variable: Stress means and method N =30, Age = 18-25 years, Experimental group I (Autogenic training), Experimental group II (meditation), Control group III (No training committed), Training duration = 6 week, Experimental design = pre and post test random group design, Statistics prouder ANCOVA Level of significance = 0.05% and Questionnaire stress through (PSS) by Everly and Girdano. Within the limitations and delimitations set for the present study and considering the results obtained, the following conclusions were drawn: The stress level was significantly decreased of all groups due to six weeks of autogenic training and meditation in judo players in compression to control group. Meditation was much effective than autogenic training in controlling the stress level in judo players.

INTRODUCTION

Many people are unhappy in the world, dissatisfied and yet not sure what is lacking in their lives. The basic reason for this unhappiness is our attachment to the material plane of existence. Once we gain a glimpse of higher spheres of consciousness, then our unhappiness and discontent automatically fade away. Man has made tremendous advances in almost every walk of the life. Modern scientists and researchers have absolutely changed his lifestyle. Yet, he is still fighting to know himself. (M.L.Gharote1999)

VARIABLES

INDIPENDENT VARIABLE: Autogenic Training and Meditation

DEPENDENT VARIABLE: Stress

MEANS AND METHODOLOGY

1. N =30
2. Age = 18-25 years
3. Experimental group I (Autogenic training)
4. Experimental group II (meditation)
5. Control group III (No training committed)
6. Training duration = 6 week
7. Experimental design = pre and post test random group design
8. Statistics = ANCOVA
9. Level of significance = 0.05%
10. Questionnaire = stress through (PSS) by Everly and Girdano

**THE TRAINING SCHEDULE ADOPTED FOR
 THE RESEARCH WORK**

Control group	Experiment group I	Experiment group II
NO EXPERIMENTAL TREATMENT	EXPERIMENTAL TREATMENT 1. AUTOGENIC TRAINING	EXPERIMENTAL TREATMENT 1. MEDITATION
	TRAINING THREE DAYS A WEEK (MONDAY, WEDNESDAY, FRIDAY) Duration :- 30 min /day	TRAINING THREE DAYS A WEEK (TUESDAY, THURSDAY, SATURDAY) Duration :- 30 min /day

**ANALYSIS OF COVARIANCE OF THE MEANS OF TWO EXPERIMENTAL GROUPS
 AND THE CONTROL GROUP IN STRESS**


Mean	C.Gr	A.T. Gr.	M. Gr.	S.V.	d.f.	S.S.	M.S.	Obt. F	Tab. F
Pre-test	25	23.3	24.8	$\frac{B/W}{W/N}$	$\frac{2}{27}$	$\frac{17.26}{699.7}$	$\frac{8.63}{25.91}$	0.33	3.35
Post test	24.8	23.2	19.6	$\frac{B/W}{W/N}$	$\frac{2}{27}$	$\frac{141.8}{643.6}$	$\frac{70.93}{23.83}$	2.97	3.35
Adjusted post mean	10.4	14.3	20.07	$\frac{B/W}{W/N}$	$\frac{2}{26}$	$\frac{80.23}{217.8}$ 4	$\frac{40.11}{8.37}$	4.78*	3.37

Significant Table F ratio at 0.05 level of confidence for 2 and 27 (df) =3.35, 2 and 26 (df) =3.37C.G.- control group, A.T.-Autogenic Training, M. – Meditation, S.V.- Source of variance,

**TURKEY POST HOC COMPARISON OF TWO EXPERIMENTAL AND CONTROL
 GROUPS IN RELATION TO STRESS**

Control group	Autogenic training	Mean deviation	Mean difference	Confidential interval value
10.4	14.9		3.79*	3.19
10.4		20.7	9.67*	
	14.9	20.7	5.88*	

**GRAPH ON ORDER TO ADJUSTED MEAN OF PHYCHOLOGICAL VARIABLE
 STRESS**



RESULTS OF STRESS

Table shows that the pre test means of control group, autogenic training and meditation were 25, 23.3, and 24.8 respectively. The table F ratio value was 3.35. When compared to table F value the obtained F value of 0.33 was lower and not significant at 0.05 level of confidence for the degrees of freedom 2 and 27. The post test means of control group, autogenic training and meditation were 24.8, 23.2, and 19.6 respectively. The table F ratio value was 3.35. When compared to table F value the obtained F value of 2.98 was lower and not significant at 0.05 level of confidence for the degrees of freedom 2 and 27.

The adjusted post test means of control group, autogenic training and meditation were 10.40, 14.19, and 20.07 respectively. The table F ratio value was 3.37. When compared to table F value the obtained F value of 4.79 was greater and significant at 0.05 level of confidence for the degrees of freedom 2 and 26. The adjusted post test means of three groups. The adjusted post test means for control, autogenic training, and meditation were 10.4, 14.9, and 20.07 respectively. The mean difference between control and autogenic training groups, control and meditation groups and autogenic training and meditation group were 3.79, 9.67, and 5.88 respectively. The Turkey's confidence interval value was 3.19.

DISCUSSION ON FINDINGS OF STRESS

Findings of the study shown that there was a significant decrease in stress level of judo players ,due to SIX WEEKS TRAINING OF autogenic training and meditation

CONCLUSIONS

Within the limitations and delimitations set for the present study and considering the results obtained, the following conclusions were drawn: The stress level was significantly decreased of all groups due to six weeks of autogenic training and meditation in judo players in compression to control group. Meditation was much effective than autogenic training in controlling the stress level in judo players.

RECOMMENDATIONS

The following recommendations have been derived on the basis of this study. The present study shows that there was a significant decrease in the stress level due to influence of six week autogenic training and meditation. Hence autogenic training meditation could be included as one of the relaxation methods to reduce the level of mental stress in judo players. Further, it is recommended that autogenic training and meditation could be used as the relaxation methods to reduce anxiety and stress in judo players.

References

- Agarwal (1988). *Research in Emerging Fields of Education Concepts, Trends and Prospects*. Sterling Publisher Private Limited, New Delhi, p. 35.
- American Association of Health, Physical Education and Recreation (1992). *Youth Fitness Test Manual*. AAHPER, Washington D.C.
- Barrow and McGee (1982). *A Practical Approach to Measurement in Physical Education*. p.112.
- Dennis Child (1972). *The Essentials of Factor Analysis*, Holt, Rinehart and Winton Publishers, London, p.45.
- Fleishman Edvin, A (1963). *The Structure and Measurement of Physical Fitness*, Englewood Cliffs, Prentice Hall Inc., New Jersey.
- Guliford, J P (1978). *Fundamental Statistics in Psychology and Education*, McGraw Hill International Publishers, Singapore.
- Hardyal Singh (1991). *Science of Sports Training*. DVS Publications, New Delhi, p. 15.